



Estimation of Arsenic and Mercury in a Polyherbal Formulation - Septiloc

Naithani V*, Pathak N, Chaudhary M

Intellectual Scientific Division, Venus Medicine Research Centre, Hill Top Industrial Estate, Bhatoli Kalan, Baddi, H.P. - 173205 India

ABSTRACT

Herbal preparations are getting popular day by day as they are considered to be safe. Septiloc is also a herbal preparation which is useful in a wide range of wounds such as diabetic wounds, bed sores, ulcers, burns, post-surgical wounds etc. It was subjected to analysis by AAS-VGA for toxic heavy metals like arsenic and mercury. Levels of arsenic and mercury were estimated in ten different batches of Septiloc and evaluated with respect to permissible limit (PL) and provisional maximum tolerable daily intake (PMTDI). In most of the samples, As and Hg were either below the detection limit or below the PL and PMTDI. These results prove that Septiloc is a safe polyherbal preparation and is free from toxic effects of As and Hg.

Keywords: Arsenic, Mercury, Septiloc, AAS-VGA.

INTRODUCTION

Arsenic and mercury are considered to be the most toxic volatile heavy metals. Arsenic is reported to cause hypertension, peripheral arteriosclerosis, skin diseases and neurotoxicity^[1-4] whereas mercury causes neurological disorders, paralysis, digestive-tract inflammation, uremia, acrodynia and immunotoxicity.^[5-9]

Earlier researchers have reported the presence of As and Hg in various herbal preparations such as in ginseng products^[10]; natural drugs^[11]; traditional Chinese medicines^[12]; Dashmoola^[13]; mercury in *Smilax myosotiflora* herbal preparations and tongkat Ali hitam herbal preparations^[14]; arsenic poisoning from anti-asthmatic herbal preparations^[15] etc. These reports throw light on the fact that medicinal plants which are considered harmless and are used as starting material for any herbal product may also contain toxic metals like As and Hg. This study was done to assess and confirm the safety of Septiloc (a polyherbal preparation for non-healing chronic wounds) in relation to As and Hg levels.

MATERIAL AND METHODS

Samples of ten different batches of Septiloc (Table 1) were

***Corresponding author: Dr. Vijay Naithani**

Intellectual Scientific Division

Venus Medicine Research Centre

Hill Top Industrial Estate, Bhatoli Kalan, Baddi,

H.P. - 173205 India; **Phone No:** 91-1795-302160

Fax No: 91-172-2565566;

E-mail: vijaynaithani@venusremedies.com

procured from Venus Remedies Limited, Panchkula, India and were subjected to analysis by Atomic Absorption Spectrometer (AAS-VGA).

All the glasswares were of Borosil "A" grade. Deionised water was used throughout the study. All the chemicals used were of analytical grade (AR). Mixed working standard (1 and 10 microgram/ml) solutions were freshly prepared by diluting the stock solutions of 1000 microgram/ml (Merck India). Cold digestion for volatile heavy metals was followed and the method was developed and standardized in the laboratory. Weighed powdered sample (0.1 g) was digested in Erlenmeyer flask (100 ml) and the flask was left overnight after adding 10 ml of conc. Sulphuric acid. It was then incubated at 70°C in a water bath for one hour. The flask was then placed in an ice bath with constant shaking saturated aqueous potassium permanganate solution was added slowly. The process was continued till the colour of the permanganate persisted. After the flask reached room temperature, one ml of hydroxylamine hydrochloride (20 % w/v in distilled water) was added to reduce excess potassium permanganate. This solution was made to desired volume by deionized water and used for estimation of As and Hg by Atomic Absorption Spectrophotometer (AAS) with Vapour Generation Assembly (VGA).

RESULTS AND DISCUSSION

Levels of arsenic and mercury were estimated in ten different batches of Septiloc and evaluated with respect to permissible limits and provisional maximum tolerable daily intake. Arsenic was detected in five samples only and was found to

be highest in SEP05 (0.28±0.04 mg/kg) and lowest in SEP02 (0.05±0.01 mg/kg). The US-FDA's allowable limits for arsenic are 0.5 ppm in uncooked muscle and 2.0 ppm in liver and other edible uncooked tissues. Results were below this limit ^[16] and Provisional maximum tolerable daily Intake (PMTDI) ^[17-20] of 0.10 mg/kg body weight in all the samples. Mercury was detected in seven samples only and was found to be highest in SEP05 (0.51±0.06 mg/kg) and lowest in SEP09 (0.10±0.02 mg/kg). It was below the permissible limit of 1.0 mg/Kg ^[21] and Provisional maximum tolerable daily Intake (PMTDI) ^[17-20] of 0.30 mg/kg body weight in all the samples. Mean levels were also within the safe limits (0.07 mg/kg for As and 0.18 mg/kg for Hg) and were found to be below the permissible limits and Provisional Maximum Tolerable Daily Intake (PMTDI). These results prove that Septiloc is a safe polyherbal preparation as far as As and Hg contamination is concerned.

Table 1 : As and Hg content in Septiloc

Sample Code	As (mg/kg)	Hg (mg/kg)
SEP01	0.09±0.02	0.21±0.03
SEP02	0.05±0.01	0.12±0.02
SEP03	BDL	0.30±0.04
SEP04	BDL	BDL
SEP05	0.28±0.04	0.51±0.06
SEP06	0.20±0.03	0.43±0.03
SEP07	BDL	BDL
SEP08	0.11±0.02	BDL
SEP09	BDL	0.10±0.02
SEP10	BDL	0.13±0.01

BDL : Below Detection Limit
Values are Mean±S.D. of 3 readings each.

REFERENCE

1. Lee YL, Shih MC, Wu WJ, Chou YH, Huang CH. Clinical and urographic presentation of transitional cell carcinoma of the ureter in a Blackfoot disease endemic area in Southern Taiwan. *Kaohsiung J Med Sci* 2002; 18: 443-449.
2. Lee MY, Jung BI, Chung SM, Bae ON, Lee JY, Park JD, Yang JS, Lee H and Chung JH. Arsenic-induced dysfunction in relaxation of blood vessels. *Environ Health Perspect* 2003; 11: 513-517.
3. Cabrera HN, Gomez MI. Skin cancer induced by arsenic in water. *J Cutan Med Surg* 2003; 7: 106-111.
4. Vahidnia A, van der Voet GB, de Wolff FA. Arsenic neurotoxicity-A review. *Human and Experimental Toxicology* 2007; 26: 823-832.

5. Langford NJ, Ferner RE. Toxicity of mercury. *Journal of Human Hypertension* 1999; 13: 651-656.
6. ATSDR. Toxicological Profile for Mercury: Agency for Toxic Substances and Disease Registry, Atlanta, GA, 1999.
7. Sweet LI, Zelikoff JT. Toxicology and immunotoxicology of mercury: a comparative review in fish and humans. *J Toxicol Environ Health B Crit rev* 2001; 4: 161-205.
8. TCE. Mercury Poisoning in: *The Columbia Encyclopedia*, sixth edition, Colombian university Press, Columbia, 2008.
9. Celebi N, Canbay O, Aycan IO, Sahin A, Aypar U. Mercury intoxication and neuropathic pain. *Paediatr Anaesth* 2008; 18: 440-442.
10. Khan IA, Allgood J, Walker LA, Abourashed EA, Schlenk D, Benson WH. Determination of heavy metals and pesticides in ginseng products. *Journal of AOAC* 2001; 84: 936-939.
11. Chuang IC, Chen KS, Huang YL, Lee PN, Lin TH. Determination of trace elements in some natural drugs by atomic absorption spectrometry. *Boil Trace Elem Res* 2000; 76: 235-244.
12. Wen HM, Chen XH, Dong TX, Zhan HQ, Bi KS. Determination of heavy metals in four traditional Chinese medicines by ICP-MS. *Zhongguo Zhong Yao Za Zhi* 2006; 31: 1314-1317.
13. Rai V, Kakkar P, Singh J, Misra C, Kumar S, Mehrotra S. Toxic metals and organochlorine pesticides residue in single herbal drugs used in important ayurvedic formulation-Dashmoola. *Environmental Monitoring Assessment* 2008; 143: 273-277.
14. Ang HH, Lee KL. Evaluation of mercury contamination in *Smilax myosotiflora* herbal preparations. *International Journal of Toxicology* 2007; 26: 433-439.
15. Tay CH, Seah CS. Arsenic poisoning from anti-asthmatic herbal preparations. *Med J Aust* 1975; 2: 424-428.
16. FDA. U.S. Food and Drug Administration, U.S. Department of Health and Human Services, 21 CFR 556.60, 1999.
17. World Health organization. Evaluation of certain food additives and contaminants, Thirty-third report of the joint FAO/WHO expert committee on Food Additives. WHO Technical Report Series number 776. WHO, Geneva, Switzerland, 1989.
18. World Health Organization. Guideline for drinking water quality. Second edition, vol.1, Recommendations. WHO, Geneva, Switzerland, 1997.
19. World Health Organization. Quality control methods for medicinal plant materials. WHO, Geneva, Switzerland, 1998.
20. JECFA. Evaluation of certain food additives and contaminants. Forty-first report of the joint FAO/WHO expert committee on food additives, Geneva, Switzerland, 1993.
21. FDA. US Food and Drug Administration: Action Level for Methyl mercury in Fish, Federal Register 49, Nov. 19, 45663, 1984.