



Short Communication

ISSN: 0975-248X
CODEN (USA): IJPSPP

**Biologically Active Triterpenes from *Eugenia pisiformis* Cambess.
(Myrtaceae)**

C. A. Oliveira, C. B. Viana, R. C. C. Martins*

Federal University of Rio de Janeiro, Institute of Natural Products Research, 21941-902, Rio de Janeiro, RJ, Brazil

ABSTRACT

In Brazil, *Eugenia pisiformis*, which belongs to botanical family Myrtaceae occurs in the states of Rio de Janeiro, Espírito Santo, Maranhão and Bahia. A specimen of *E. pisiformis* collected in Itatiaia National Park, State of Rio de Janeiro had its leaves, were studied because there are no phytochemical data in the literature for this plant so far. From the ethanolic extract of the leaves 3 substances were isolated and identified as the cycloartane-type triterpene 24-methylenepollinastanol, oleanane-type triterpene uvaol and ursane-type triterpene erythrodiol. All of them present significant biological properties, according to the literature and 24-methylenepollinastanol have never been isolated from this species, genus or family.

Keywords: Myrtaceae, *Eugenia pisiformis*, triterpene, cycloartane.

DOI: 10.25004/IJPSDR.2017.090211

Int. J. Pharm. Sci. Drug Res. 2017; 9(2): 98-99

INTRODUCTION

Myrtaceae is a plant family, which comprises about 142 genera, 17 tribes and 5500 species distributed in temperate tropical regions in both hemispheres. [1-2] *Eugenia* is a genus from Myrtaceae that comprises about 1,009 species, spreading from Mexico to Argentina. In Brazil, it occurs mainly in the Midwest and Southeast regions, as trees or shrubs. From *Eugenia* species have been isolated several types of secondary metabolites, mainly phenolic, including flavonoids, catechins and tannins, as well as terpenoid substances [3-5] many of them possessing significant biological activities such as anti-inflammatory and bactericidal properties. *Eugenia pisiformis* Cambess. is a species that

occurs in the Atlantic Forest, mainly in states of Rio de Janeiro, Espírito Santo, Maranhão, Paraná and Bahia. As a part of a study which aims to search biologically active compounds in the genus Myrtaceae, *E. pisiformis* was chosen because there are, so far, no records of phytochemical studies for this plant in the literature.




Fig. 1: Triterpenes 24-methylenepollinastanol (1), uvaol (2) and erythrodiol (3).

***Corresponding author: Dr. R. C. C. Martins,**

Adjunct Professor, Laboratório de Pesquisa de Metabolismo Especial, Institute of Natural Products Research, HSS-20, 21941-902, Rio de Janeiro, RJ, Brazil;
E-mail: roberto@nppn.ufrj.br

Received: 23 February, 2017; **Accepted:** 15 March, 2017

MATERIALS AND METHODS

Leaves of *E. pisiformis* were collected at Itatiaia National Park, in Rio de Janeiro state, Brazil in May, 2014. Fresh material was dried in oven at 40°C during 48 hours and grounded to a fine powder in a knife mill. Powder of the leaves was then extracted by percolation with cold

ethanol and the resultant ethanolic extract was submitted to several chromatographic steps. These procedures led to the isolation and further identification of 3 triterpenes by HRESI-MS and NMR.

RESULTS AND DISCUSSION

Cycloartane-type *nor*-triterpene 24-methylenepollinastanol ((3 β , 5 α)-14-methyl-9, 19-cycloergost-24(28)-en-3-ol) (**I**) was isolated and structurally characterized through analysis by 1D and 2D NMR spectra. Molecular formula C₂₉H₄₈O was confirmed by HRESI-MS experiments. Ursane-type triterpene uvaol (3 β -urs-12-ene-3, 28-diol) (**II**) and oleanane-type erythrodiol (3 β -olean-12-ene-3, 28-diol) (**III**) were also isolated and structurally characterized by 1D and 2D NMR and HRESI-MS (Fig. 1).

24-methylenepollinastanol has already been isolated from the following species: *Holarrhena curtisii* (Apocynaceae) [6], *Musa sapientum* (Musaceae) [7], *Cajanus cajan* (Fabaceae) [8], Fenugreek (Fabaceae) [9], Dwarf cavendish (Musaceae). [10] Although there is not yet any biological property related to this compound, literature describes cycloartane triterpenes as biologically active substances, showing important properties such as cytotoxic [11], antiprotozoal [11], antiacetylcholinesterase [12] and anti-inflammatory [13] among several others. Thus, 24 methylenepollinastanol might be a potential candidate as a biologically active substance. Additionally, an extensive search in the literature and virtual databases confirmed that this compound have never been isolated before from this species, from the genus *Eugenia* or Myrtaceae. This class of triterpenes is very uncommon in both *Eugenia* and Myrtaceae.

Triterpene uvaol has already been isolated from several plants used in folk medicine and it has confirmed antioxidant [14], antibacterial [15], antiviral [16] and anti-inflammatory [17] activities. In the genus *Eugenia*, there are reports of the triterpenes uvaol and erythrodiol in the species: *E. umbelliflora*, *E. uniflora* and *E. catharinae*. [18, 4]

As a first phytochemical study, *E. pisiformis* has showed itself as a potential source of biologically active triterpenes. It has also showed to be able to metabolize uncommon cycloartane-type triterpenes. Phytochemical studies keep on going, as these are only the preliminary results for this plant.

ACKNOWLEDGEMENTS

Authors would like to thank CAPES for the scholarship and IPPN-UFRJ for the financial support for this work.

REFERENCES

- Retamales HA, Scharaschkin T. Comparative leaf anatomy and micromorphology of the Chilean Myrtaceae: Taxonomic and ecological implications. *Flora*. 2015;217:138-154.
- Wilson PG, O'Brien MM, Heslewood MM, Quinn CJ. Relationships within Myrtaceae sensu lato based on a matk phylogeny. *Plant Syst and Evol*. 2005; 251:3-19.
- Amorim ACL, Lima CKF, Hovell AMC, Miranda ALP, Rezende CM. Antinociceptive and hypothermic evaluation of the leaf essential oil and isolated terpenoids from *Eugenia uniflora* L. (Brazilian Pitanga). *Phytomedicine*. 2009; 16:923-928.
- Moresco HH, Colla G, Cavalcante ID, Queiroz G S, Pizzolatti MG, Brighente IM. Chemical constituents of *Eugenia catharinae* and their antioxidant activity. *Natural Product Research*. 2016; 13:1-5.
- Sant'Ana AEG, Oliveira AM, Humberto MMS, Silva JM, Rocha RFA. Estudo fitoquímico e avaliação das atividades moluscicida e larvicida dos extratos da casca do caule e folha de *Eugenia malaccensis* L. (Myrtaceae). *Revista Brasileira de Farmacognosia*. 2006; 16:618-629.
- Pornpakakul S, Srisurichan S. Triterpenoids from the seedpods of *Holarrhena curtisii* King and Gamble. *Phytochem Lett*. 2015; 12:282-286.
- Knapp FF, Phillips DO, Goad LJ, Goodwin TW. Isolation of 14 α -methyl-9 β ,19-cyclo-5 α -ergost-24(28)-en-3 β -ol from *Musa sapientum*. *Phytochemistry*. 1972; 11: 3497- 5000.
- Akihisa T, Nishimura Y, Nakamura N, Roy K, Ghosh P, Thakur S.; Tamura T. Sterols of *Cajanus cajan* and three other leguminosae seeds. *Phytomedicine*, 1992; 31(5): 1765-1768.
- Cerdon C, Rahier A, Taton M, Sauvaire Y. Effect of tridemorph and fenpropimorph on sterol composition in *Fenugreek*. *Phytochemistry*. 1996; 41(2): 423-431.
- Oliveira L, Freire CSR, Silvestre AJD, Cordeiro N, Torres IC, Evtuguin D. Lipophilic extractives from different morphological parts of banana plant "Dwarf Cavendish". *Industrial Crops and Products*. 2006; 2: 201-211.
- Morais SM, Silva AAS, Falcão MJC, Vieira IGP, Ribeiro LM, Viana SM, Teixeira MJ, Barreto FS, Carvalho CA, Cardoso RPA, Andrade-Junior HF. Activity of cycloartane-type triterpenes and sterols isolated from *Musa paradisiaca* fruit peel against *Leishmania infantum chagasi*. *Phytomedicine*. 2014; 21:1419-1423.
- Zahid S, Ata A, Samarasekera, R. New Cycloartane-type Triterpenoids from *Artocarpus nobilis*. *Z. Naturforsch*, 2007, 62: 280-284.
- Romero JC, Martínez-Vázquez A, Herrera MP, Martínez-Mayorga K, Parra-Delgado H, Pérez-Flores FJ, Martínez-Vázquez, M. Synthesis, anti-inflammatory activity and modeling studies of cycloartane-type terpenes derivatives isolated from *Parthenium argentatum*. *Bioorg Med Chem*. 2014; 22: 6893-6898.
- Allouche Y, Warleta F, Campos, M, Sanchez-Quesada C, Uceda M, Beltran G; Gaforio JJ. Antioxidant, antiproliferative, and pro-apoptotic capacities of pentacyclic triterpenes found in the skin of olives on MCF-7 human breast cancer cells and their effects on DNA damage. *J Agric and Food Chem*. 2011; 59:121-130.
- Martins A, Vasas A, Viveiros M, Molnar J, Hohmann J, Amaral L. Antibacterial properties of compounds isolated from *Carpobrotus edulis*. *Int Journal of Antimicrobial Ag*. 2010; 37: 438-444.
- Serra C, Lampis G, Pompei R, Pinza M. Antiviral activity of new triterpenic derivatives. *Pharmacol Res*. 1994; 29:359-366.
- Barreto E, Agra LC, Lins MP, Marques OS, Smaniotto S, Melo CB, Lagente V. Uvaol attenuates pleuritis and eosinophilic inflammation in ovalbumin-induced allergy in mice. *Eur Journal of Pharmacol*. 2016; 780:232-242.
- Nogueira JMR, Queiroz JMG, Suzuki MCM, Motta APR, Carvalho EM. Aspectos populares e científicos do uso de espécies de eugenia como fitoterápico. *Rev Fitos*. 2015; 9: 73-159.

Source of Support: Nil, Conflict of Interest: None declared.